

DETECTION OF ANTI-CORRELATED HARD X-RAY TIME LAG IN CYGNUS X-3

MANOJENDU CHOUDHURY¹ & A. R. RAO
Tata Institute of Fundamental Research, Mumbai-400005, India

Received 2004 August 17; Accepted 2004 October 07

ABSTRACT

The wide-band X-ray spectra of the high mass X-ray binary Cygnus X-3 exhibits a pivoting behavior in the ‘low’ (as well as ‘hard’) state, correlated to the radio emission. The time scale of the soft and hard X-rays’ anti-correlation, which gave rise to the pivoting feature, was found to be less than a day from the monitoring observations by *RXTE*–ASM and *CGRO*–BATSE. In this *Letter* we report the detection of a lag of $\lesssim 1000$ s in the anti-correlation of the hard X-ray emission (20–50 keV) to that of the soft X-ray emission (2–7 keV), which may be attributed to the viscous time scale of flow of matter in the accretion disk. This suggests the geometrical picture of a truncated accretion disc with a Compton cloud inside the disc, the relative sizes of which determine the spectral shape. Any change in the disc structure will take place in a viscous time scale, with corresponding anti-correlated change in the Compton cloud. We also report the pivoting in the spectra in one span of a pointed observation when an episode of the rearranging of the accretion system is serendipitously observed. This is the first such observation of hard X-ray delay seen in the persistent Galactic microquasars, within the precincts of the hard state.

Subject headings: accretion – binaries : close – stars : individual (Cygnus X-3) – X-rays : binaries

1. INTRODUCTION

The presence of a non-thermal component in the wide-band X-ray spectrum is one of the most prominent observational features of the Galactic blackhole candidates as well as neutron star binaries (see Barret 2004, for a short review), although the very precise nature of its origin has not yet been definitively ascertained. Conflicting views regarding the physical mechanism of the generation of hard X-ray emission, ranging from synchrotron at the base of the jet (Markoff et al. 2001, 2003) to Comptonization of thermal photons by a hot corona with various geometrical structures (see, for eg., Poutanen 1998) – the most favored being that of a hot quasi-spherical cloud inside a truncated disc (Zdziarski et al. 2002) – exist in the literature. McClintock & Remillard (2004) establish the various classes of X-ray emission states, based on the spectral characteristics, adding to the canonical ‘low-hard’ and ‘high-soft’ states (Tanaka & Lewin 1995). Of these, the ‘low-hard’ state is perhaps the most interesting for all blackhole sources for the presence of ubiquitous outflow in the form of a collimated jet (Fender 2001). These jets are also present in the low magnetic field ($\lesssim 10^9$ G) neutron stars (Fender & Kuulkers 2001), collectively these objects are classified as *microquasars* (Mirabel & Rodriguez 1999). The ADAF (Narayan & Yi 1994) and TCAF (Chakrabarti 1996) models provide viable alternative explanations of the accretion-ejection phenomena (Meier 2001; Das & Chakrabarti 1999, respectively) in this class of sources, involving a truncated accretion disc.

Cygnus X-3 is an enigmatic X-ray binary system which is one of the brightest Galactic sources in the radio band. The X-ray (as well as the infra-red) emission exhibit a binary modulation of 4.8 hours (Singh et al. 2002). Al-

though the nature of the compact object is not yet known, the persistent radio emission from a jet classifies it as a prominent microquasar, and the different X-ray spectral states of this source (Choudhury & Rao 2002) are similar in nature to that of the canonical blackhole candidates, although individual components describing the spectral shape may be different (Choudhury et al. 2003). The long term statistical correlation among the soft X-ray (2–12 keV, *RXTE*–ASM), hard X-ray (20–100 keV, *CGRO*–BATSE) and radio emission (2.2 GHz, *GBI*) revealed a pivoting behavior of the X-ray spectra (in the region of 10–20 keV) correlated to the radio emission, within the precincts of the hard state of this source (Choudhury et al. 2002). This behavioral feature, in the ‘low-hard’ state, has been established as a general feature of the X-ray emission for the Galactic blackhole binaries, with the pivot point varying for the various sources (Choudhury et al. 2003). In Choudhury et al. (2004) it has been established, from the daily monitoring of the source simultaneously in the soft X-ray, hard X-ray and radio, that the correlation time scale for Cygnus X-3 is less than a day. Considering the universal correlation of the soft X-ray flux with the radio spanning 5 orders of magnitude of intrinsic luminosity, Choudhury et al. (2003) postulated that it is the soft X-ray that determines the spectral pivoting phenomena which is the driving force of the evolution of the accretion-ejection mechanism in these systems. In such a scenario, the Compton scattered X-rays originate from a region close to the compact object, which may be confined within the Centrifugal Boundary Layer (CENBOL) (Chakrabarti & Titarchuk 1995; Ebisawa et al. 1996). At low accretion rates, the disk is truncated far away from the compact object and the X-ray spectrum is dominated by a thermal-Compton spectrum, originating from the high

¹Current address: N.C.R.A., T.I.F.R., Pune-411007. India.
e-mail: manojendu@ncra.tifr.res.in

temperature region within. Increase/decrease in accretion rate causes the location of the disk truncation (CEN-BOL) to move in/out, causing the Comptonizing cloud to decrease/increase as well as the amount of seed thermal photon to increase/decrease which cools the Comptonizing cloud more/less efficiently, giving rise to the pivoting of the wide-band X-ray spectrum. Hence, the pivoting of the X-ray spectrum is not expected to occur instantaneously, as the viscous time scale of the flow of matter in the accretion disc determines the change in the hard X-rays vis-a-vis the soft X-rays, and in such a situation the delay of the hard X-ray should be much less than a day (Choudhury et al. 2004), implying the need to search for delay between the hard and soft X-rays using the long pointed observations, spanning at least a good fraction of one binary orbit. Considering the fact that the mechanism of a change in accretion rate is very unlikely to be a continuous process, the individual events of such a change in the soft X-ray emission with an eventual anti-correlated change in the hard X-ray emission need to be serendipitously observed during the pointed observations, with the source in the ‘low’ (as well as ‘hard’) state.




Figure 1. A typical cross-correlation between the soft (2-7 keV) and hard (20-50 keV) X-ray flux with the lightcurve not corrected for binary modulation. This particular result was obtained for observations on MJD 50953.

In this *Letter*, we report the detection of such an anti-correlated delay, of $\lesssim 1000$ s of the hard X-rays (20–50 keV) with respect to the soft X-ray flux (2–7 keV), i.e. any change in the soft X-ray flux results in an opposite change in the hard X-ray flux after the observed delay time scale, during the individual pointed observations using the PCA and HEXTE aboard the *RXTE*, in the ‘low’ (correspondingly ‘hard’) state of the X-ray emission in the source. This type of lagged anti-correlation, we believe, is the first to be observed for the class of X-ray persistent Galactic microquasars within the bounds of the (low-) hard state. We confirm our detection of this anti-correlated delay by observing the pivoting of the spectra ‘in-situ’ during one of the pointed observations of the *RXTE*-PCA.

2. DATA AND ANALYSIS

The data are obtained from the archives of the *RXTE* through the HEASARC Online Service. The pointed observations of both the narrow field of view instruments aboard the *RXTE*, viz. PCA and HEXTE (cluster 0) are used for initial study of the cross-correlation of the lightcurves at the various energy bands, whereas the results of binary corrected lightcurves are reported here only for the PCA observations. The lightcurves as well as the spectrum (obtained from the PCA) uses the standard 2

form of data (all PCUs added), with all the procedures of data filtering, background and deadtime corrections strictly adhered to. The basic data reduction and analysis was carried out using HEASOFT (v5.2). The correction for binary modulation was done using the recipe described by Choudhury et al. (2004), using the binary template of van der Klis & Bonnet-Bidaud (1989) and obtaining the phase of each pointing from the quadratic ephemeris of Singh et al. (2002).




Figure 2. The (binary uncorrected) lightcurve of the source on MJD 50321 as observed by the PCA at 2-7 keV (top panel) and 20-50 keV (middle panel), along with the HEXTE at 20-60 keV (bottom panel). The cross-correlation shows a clear indication of possible anti-correlated delay of the hard X-ray flux.

The pointed observations of this source from the *RXTE* were done predominantly in the ‘high’ state, with the emphasis on the investigation during the massive radio flaring episodes. As a consequence, only a few observations in the ‘low’ state exist for this source and that too during only two phases, MJD 50319–50325 and 50949–50954. During the first phase the electron background was high and one needs to filter the data very judiciously, while on the other occasion the source exhibited its ‘hardest’ state, providing one extreme of the X-ray spectral characteristics. The binary modulation of the X-ray emission is very strong in the source. In the ‘low’ state, the soft X-ray flux (2–7 keV) is modulated by a factor ~ 2 and the hard X-ray flux (20–50 keV) is modulated by a factor ~ 2.5 –3. This strong modulation smothers all other variations in nearly all the cases, and hence the cross-correlation shows a strong correlation between the emission at the two flux ranges (Figure 1). Extensive search for anti-correlated delay from the binary modulated lightcurves yielded only one possible indication of the presence of such a delay, on MJD 50321 (obs. id.: 10126-01-01-01). The strong anti-correlation with a delay of ~ 500 –700 s is seen for hard X-rays from both PCA (20–50 keV) and HEXTE (20–60 keV) with respect to the soft X-rays (2–7 keV, PCA), as illustrated in Figure 2. The positive correlation without any delay, due to the binary

modulation is as prominent (F.F.T. correlation coefficient - hereafter CC - of 0.3 for 488 degrees of freedom - hereafter dof, Pearson's CC of 0.56 for 244 dof - with a null probability ~ 0 , and Spearman's CC of 0.54 with 488 dof — null probability ~ 0) as the negative correlation with a delay of a few hundred seconds (CC of -0.6 for the PCA data). When the smooth and strong binary variation is subtracted using the binary ephemeris (Choudhury et al. 2004), the positive correlation got suppressed and the negative correlation with a delay was the only prominent feature of the cross-correlation plot.




Figure 3. The lightcurve as observed by the PCA at 2-7 keV and 20-50 keV, both uncorrected and corrected for the binary modulation, on the five days showing the anti-correlated hard X-ray lags. The cross-correlation plots shown are obtained after co-adding the lightcurves of MJD 50321 & 50322, and MJD 50952, 50953 & 50954 (two phases), respectively, to compute the maximum possible error in the measurement. The details of the individual pointed observations are given in table 1.

Repeating the cross-correlation test for these ‘low’ (and ‘hard’) states after correcting for the binary modulation (Choudhury et al. 2004) provided very significant results, showing a very strong anti-correlated delay of the hard X-rays (20–50 keV, PCA) with that of the soft X-rays (2–7 keV, PCA), on five days, including MJD 50321. Table 1 gives the details of the days of observation of the delay, the observation id and the time delay observed on the particular pointings. The delay is first obtained by the F.F.T. algorithm using the XRONOS program “crosscor”, the Pearson’s and Spearman’s coefficients are subsequently obtained. The F.F.T. coefficient is normalized by (dividing by) the square root of the product of number of good newbins of the concerned lightcurves, effectively this coefficient is the cross covariance of the two lightcurves. To substantiate the statistical robustness of the observed delay we co-add the lightcurves (in two phases, MJD 50321-22 & 50952-54) to obtain the maximum possible error in the correlation coefficient. The cross-correlation of the co-

added lightcurves, along with the individual lightcurves (both binary uncorrected and corrected) are presented in Figure 3, where the delay as well as cross-correlation is established, despite the intrinsic variation in the delay as well as the correlation coefficient of the individual pointings (table 1).

The spectral evolution of the source during the pointed observation on MJD 50321 (obs. id.: 10126-01-01-01) is shown in Figure 4, where the two spectra correspond to the softer and harder regions of the lightcurve, i.e. before and after pivoting. The pivoting of the spectra, earlier reported for observations taking place on different days corresponding to the extreme cases within the bounds of the ‘low’ and ‘hard’ states (Choudhury et al. 2002, 2003), is observed directly within the span of one pointed observation containing the episode of the soft X-ray flux driving the spectral evolution, and the hard X-ray flux changing correspondingly after a time scale of the viscous flow in the accreting fluid. This ‘in situ’ observation of pivoting, in the hard state, validates the cross-correlation results and puts the particular observational feature on very firm footing.

3. DISCUSSION

The strong binary modulation of Cygnus X-3 has prevented any serious study on the variability of this source at times-scales of few hundred seconds to about a day. Singh et al. (2002) have shown that the binary period (and the template of its variation) has remained consistent for more than 25 years with a linear decay with no second derivative of the period. Choudhury et al. (2004) have used this information and shown that this ephemeris can be used to remove the large amplitude smooth binary variation without recourse to any derivation of phase during individual observations - the only free parameters being the amplitude of variation and the mean count rate. Using this method, we have corrected for the large amplitude binary variations and detected anti-correlated hard X-ray delays in five of the observations.

The only assumption incorporated in our method is that the occasional jitters seen in the binary modulation is not due to any binary instability (Choudhury et al. 2004), rather we explore whether any residual energy dependent phase jitter and/or variable energy dependent template could have caused the anti-correlated delay presented here. In all observations we find a strong positive correlation with zero phase indicating the similarity of the template in the two energy bands. Further, more than 90% of the binary variability in Cygnus X-3 is energy independent (see Rajeev et al. 1994) and the large amplitude variations in high phase are correlated in all energies. The anti-correlated behavior is seen in the slow varying components which is unrelated to the binary variation. This was evident when we examined the variation of the hardness ratio compared to the lightcurves.

The strongest evidence for the reality of the anti-correlation comes from the model independent spectral pivoting seen in one of the observations (see Fig 4). Choudhury et al. (2003, 2002) have shown that this type of spectral pivoting at 10–20 keV energies occurs when one takes spectrum at different days at different count rates. The fact that such spectral pivoting is indeed seen in one single observation shows that the soft and hard X-ray anti-

correlation occurs in a time scale less than the binary period and the cross-correlation analysis reveals the delay in the time scale close to a thousand seconds.




Figure 4. The wide-band X-ray spectra of the source on MJD 50321, for the soft and the hard regions of the lightcurve, resulting in comparative softer and harder spectral distribution, displaying the ‘in situ’ pivoting in the region of 10–20 keV.

The three-way association between soft X-rays, hard X-rays and radio emission in Cygnus X-3 maybe explained by the geometry of truncated, optically thick, accretion disc, with a Comptonizing cloud of energetic electrons inside which may have advective (Narayan & Yi 1994) or bulk motion (Chakrabarti & Titarchuk 1995). The soft X-ray flux determines the accretion disk structure, while the Compton cloud within the truncated disc is responsible for the hard X-ray emission and it also drives the outflow (see Meier 2001; Das & Chakrabarti 1999, for contrast-

ing paradigms). The delay presented in this work signifies the disk readjustment time scale which should correspond to the location of the truncation radius. The viscous time t_{vis} can be calculated for the radiation pressure dominated optically thick accretion disc as

$$t_{vis} = 30\alpha^{-1} M^{-1/2} R^{7/2} \dot{M}^{-2} s$$

where α is the viscosity parameter in units of 0.01, M is the mass of the compact object in solar mass units, R is the radial location in the accretion disc in units of 10^7 cm, $\dot{M}$ is the mass accretion rate in units of 10^{18} g/s (see Belloni et al. 1997; Fender & Belloni 2004). Taking $\alpha = 1$, $M = 10$ and $\dot{M} = 3$ (corresponding to one fifth of the Eddington accretion rate), we get $R \sim 7$ for a viscous time scale of 1000 s. Thus, if Cyg X-3 harbors a black hole of $10 M_{\odot}$, the observed delay signifies a CENBOL location of ~ 25 Schwartzchild radius.

It should be noted that the X-ray emission shows a strong correlation with the radio emission, but the radio emission does not show any binary modulation. It would be interesting to repeat the type of work presented here with a simultaneous radio monitoring observations so that a clear handle can be obtained for the location of the radio emission.

ACKNOWLEDGMENTS

This research has made use of data obtained through the HEASARC Online Service, provided by the NASA/GSFC, in support of NASA High Energy Astrophysics Programs.

REFERENCES

- Barret, D. 2004, proc. of Plasmas in the Laboratory and in the Universe, American Institute of Physics Conference Series, 238-249 [astro-ph/0401100]
 Belloni, T., Mendez, M., King, A. R., van der Klis, M. & van Paradijs, J. 1997, ApJ, 488, L109
 Chakrabarti, S.K. 1996, Phys. Rep., 266, 229
 Chakrabarti, S. K., & Titarchuk, L. G. 1995, ApJ, 455, 623
 Choudhury, M., & Rao, A.R. 2002, JApA, 23, 39
 Choudhury, M., Rao, A.R., Vadawale, S.V., Jain, A.K. & Singh, N.S. 2004, A&A, 420, 665
 Choudhury, M., Rao, A.R., Vadawale, S.V. and Jain, A.K. 2003, ApJ, 593, 452
 Choudhury, M., Rao, A.R., Vadawale, S.V., Ishwara-Chandra, C.H. & Jain, A.K., 2002, A&A, 383, L35
 Das, T. & Chakrabarti, S.K. 1999, Class & Quant. Gravity, 16, 3879
 Ebisawa, K., Titarchuk, L. & Chakrabarti, S. 1996, PASJ, 48, 59
 Fender, R.P. & Belloni, T. 2004, ARA&A, 42, 317-364
 Fender, R.P. 2001, MNRAS, 322, 31
 Fender, R.P. & Kuulkers, E. 2001, MNRAS, 324, 923
 Markoff, S., Falcke, H. & Fender, R. 2001, A&A, 372, L25
 Markoff, S., Nowak, M., Corbel, S., et al. 2003, A&A, 397, 645
 McClintock, J.E. & Remillard, R.A. 2004, to appear in *Compact Stellar X-ray Sources*, W.H.G. Lewin and M. van der Klis eds., Cambridge University Press, preprint (astro-ph/0306213)
 Meier, D.L. 2001, ApJ, 548, L9
 Mirabel, I.F. & Rodriguez, L.F. 1999, ARA&A, 37, 409
 Narayan, R. & Yi, I. 1994, ApJ, 428, L13
 Poutanen, J. 1998, in *Theory of Black Hole Accretion Discs*, M. A. Abramowicz, G. Bjornsson & J. E. Pringle, eds., Cambridge Contemporary Astrophysics, Cambridge University Press, 100p.
 Rajeev, M.R., Chitnis, V.R., Rao, A.R., et al. 1994, ApJ, 424, 376
 Singh, N.S., Naik, S., Paul, B. et al. 2002, A&A, 392, 161
 Tanaka, Y. & Lewin, W.H.G. 1995, Cambridge Astrophys. Ser. vol. 26, X-Ray Binaries, eds. W.H.G. Lewin, J. van Paradijs & E.P.J. van den Heuvel (Cambridge Univ. Press), 126
 van der Klis, M. & Bonnet-Bidaud, J.M. 1989, A&A, 214, 203
 Zdziarski, A.A., Poutanen, J., Paciesas, W.A., et al. 2002, ApJ, 578, 357

TABLE 1

THE DETAILS OF THE POINTED OBSERVATIONS FOR WHICH THE LAGGED ANTI-CORRELATION OF THE HARD X-RAYS (20–50 keV, PCA) WITH RESPECT TO THE SOFT X-RAYS (2–7 keV, PCA) HAS BEEN FOUND, AFTER CORRECTING FOR THE BINARY MODULATION, AS PRESCRIBED BY CHOUDHURY ET AL. (2004). THE DETAILS OF THE DELAY OBSERVED IN THE INDIVIDUAL LIGHTCURVES OBSERVED ON THE PARTICULARLY MENTIONED DAYS IS GIVEN.

MJD	Observ. Id.	Delay (s) (error)	Statistical coefficient		
			F.F.T. (error)	Pearson (Null Prob.)	Spearman (Null Prob.)
50321	10126-01-01-01	$\sim 620 (\pm 70)$	$\sim -0.48 (\sim 0.04)$	$-0.51 (\sim 10^{-12})$	$-0.58 (\sim 10^{-30})$
50322	10126-01-01-02	$\sim 750 (\pm 120)$	$\sim -0.44 (\sim 0.04)$	$-0.46 (\sim 10^{-4})$	$-0.43 (\sim 10^{-13})$
50952	30082-04-04-00	$\sim 700 (\pm 50)$	$\sim -0.61 (\sim 0.08)$	$-0.66 (\sim 10^{-11})$	$-0.69 (\sim 10^{-36})$
50953	30082-04-05-00	$\sim 950 (\pm 60)$	$\sim -0.58 (\sim 0.08)$	$-0.73 (\sim 10^{-13})$	$-0.75 (\sim 10^{-39})$
50954	30082-04-06-00	$\sim 0-1000$	$\sim -0.40 (\sim 0.08)$	–	–